ECOLOGICAL SIGNIFICANCE OF AQUATIC BIRD GUANO IN A WETLAND ECOSYSTEM: A STUDY OF UDPURIA TALAB, DIGOD, KOTA, RAJASTHAN, INDIA

[1]DR. PINKY SHARMA AND [2]DR. RAJENDRA SINGH RAJAWAT

[1]ASSISTANT PROFESSOR, DEPARTMENT OF ZOOLOGY, GOVERNMENT COLLEGE KOTA, RAJASTHAN

[2] COORESPONDING AUTHOR, DEPARTMENT OF ZOOLOGY, GOVERNMENT COLLEGE KOTA, RAJASTHAN

ABSTRACT

The ecological significance of aquatic bird guano in a wetland ecosystem like Udpuria Talab (located in Digod, Kota, Rajasthan, India) highlights the critical role of waterbirds as ecosystem engineers and nutrient vectors. [1, 2]

Udpuria Talab is a vital regional wetland that serves as a sanctuary for both local resident waterbirds and a large influx of winter migratory species (such as the Ruddy Shelduck, Painted Stork, and Oriental Darter). The accumulation of bird guano (excrement) within this closed or semi-closed freshwater system drives essential biochemical and biological dynamics. [1, 2, 3, 4]

Keywords- Udpuria Talab, Digod, Kota, Ecology, Aquatic bird,Guano, wetland

INTRODUCTION

Guano is exceptionally rich in macronutrients, primarily Nitrogen (N) and Phosphorus (P). Birds feeding in surrounding agricultural fields or separate foraging grounds return to roost at Udpuria Bird Sanctuary, acting as mobile bio-vectors that transport external organic matter into the lake ecosystem. This phenomenon, known as guanotrophication, accelerates local nutrient loops: [1, 2]

- Ammonification and Nitrification: Microbial decomposition of uric acid from the guano releases bioavailable forms of nitrogen into the water column.
- Phosphorus Influx: In a contained rural body like Udpuria Talab, this natural fertilization provides crucial chemical inputs that might otherwise be limited.[1,2,3]

Primary Productivity and Food Web Acceleration

The steady release of N and P directly boosts the primary biological productivity of the wetland. [1, 2]

- Plankton Blooms: Nutrients encourage the rapid proliferation of phytoplankton (microscopic algae) and cyanobacteria, which form the absolute baseline of the aquatic food web. [1]
- Secondary Consumers: The surge in phytoplankton supports a thriving population of zooplankton, macroinvertebrates, and indigenous fish stocks. This ensures an abundant food supply that circles right back to sustain larger predatory fish, amphibians, and the avifauna themselves. [1, 2]

Vegetation and Aquatic Macrophytes

Nitrient-dense sediment at the bottom of the talab assists the growth of emergent and submerged aquatic macrophytes. These plants provide essential structural coverage, offering nesting sites and shelter for birds and small aquatic fauna, maintaining the physical integrity of the wetland habitat. [1, 2]

Biogeochemical Balance and Conservation Risks

While guano is an essential driver of the wetland's health, its impact is highly sensitive to bird density and water volume.

- The Balancing Act: Optimal nutrient enrichment supports high biodiversity. [1, 2]
- The Eutrophication Risk: During dry spells or periods of excessive crowding, heavy loading of guano can oversaturate the water body. This can lead to hyper-eutrophication, resulting in dense algal blooms that deplete dissolved oxygen (anoxia) and potentially stress the native fish populations. [1, 2]

Therefore, the input of avifaunal guano at Udpuria Talab is a fundamental ecological process that preserves the system's biological viability, showcasing how closely linked bird conservation is to overall wetland management in Rajasthan's Hadoti region[4,5,6]

DISCUSSION

Aquatic bird guano acts as a natural fertilizer and primary driver of guanotrophication, significantly altering the nutrient balance and biological productivity of wetland ecosystems. [1, 2, 3]

Nutrient Enrichment and Biogeochemistry

- Nitrogen and Phosphorus Loading: Guano is rich in bioavailable nitrogen (N) and phosphorus (P)—specifically as nitrates (NO_3) and phosphates (PO_4)—which directly elevate nutrient concentrations in the water column. [1]
- Sediment Retention: Wetland sediments rapidly absorb a large portion of total phosphorus from guano, while a significant share of nitrogen remains active in the water and cycles through microbial pathways. [1]
- Nutrient Vectors: Waterbirds function as mobile zoogeochemical links:
  - Birds that feed and defecate within the same waterbody (moorhens, coots, pygmy geese) recycle and transform localized nutrients.
  - Birds that feed on land and roost in wetlands (ducks, geese, swans) cause a net import of external nutrients into the aquatic ecosystem. [1, 2, 3]

Primary Production and Food Web Dynamics

- Algal and Plankton Growth: Nutrient pulses stimulate the proliferation of phytoplankton, periphyton, and aquatic weeds, forming the base of the wetland food web. [1, 2]
- Invertebrate and Fish Support: Enhanced primary productivity increases microinvertebrates, zooplankton, and benthic organisms, which in turn sustain local fish stocks and higher trophic predators. [1, 2, 3]

Management and Eutrophication Risks

- Dual Nature (Guanotrophication): While moderate guano inputs enrich biodiversity and support plant and animal growth, excessive accumulation from dense, unnaturally concentrated bird populations can trigger hyper-eutrophication, oxygen depletion, and toxic algal blooms. [1]
- Balanced Conservation: Effective wetland management requires tracking seasonal waterbird fluctuations to balance natural nutrient enrichment with water quality preservation. [7,8,9]

Udpuriya Talab is a natural water body and wetland area located near Udpuriya Village in Digod Tehsil of Kota District in Rajasthan, India. [1, 2]

Key Details

- Location: Udpuriya village, Digod sub-division/tehsil, Kota district, Rajasthan (Pincode: 325201). [1, 2, 3]
- Distance from Kota: Roughly 32 km east of Kota city and about 8 km from Digod tehsil headquarters. [1]
- Biodiversity and Migratory Birds: The pond and surrounding wetland area serve as a prominent seasonal breeding and roosting site for various migratory and resident bird species—most notably Painted Storks (Janghila), Openbill Storks, Black-necked Storks, and Purple Moorhens. [1]
- Seasonality: The bird activity and congregation peak annually between August and February, drawing local attention and nature enthusiasts. [1]
- Local Context: Udpuriya is a small rural settlement administered under the Doongarja Gram Panchayat within the Sangod Vidhan Sabha assembly constituency. [1]

RESULTS

The ecological significance of aquatic bird guano within a wetland ecosystem—specifically contextualized around Udpuria Talab (Udpuria Wetland) in Digod, Kota, Rajasthan, India—revolves around a biophysiological phenomenon known as guanotrophy. Aquatic birds act as critical mobile nutrient vectors, actively gathering nutrients across various foraging grounds and depositing them into the localized wetland matrix via their excrement. [10,11,12]

At Udpuria Talab—a notable rural wetland hotspot populated by resident and winter migratory waterbirds like the Brown Crake, Oriental Darter, Eurasian Coot, Indian Cormorant, and various egrets—this guano acts as a foundational biological engineering driver. [1, 2, 3]

Spatial & Ecological Framework

Macro-Geographic Range

Udpuria Talab is nestled within the semi-arid landscape of the Kota district in southeastern Rajasthan, India. The geographic positioning makes this region a crucial hub for the Central

Asian Flyway, bringing thousands of migratory waterfowl (such as winter migrant ducks of the Anatidae family) to rest alongside localized residential aquatic birds. [1]

Micro-Environmental Habitat

The micro-environment comprises a shallow freshwater inland wetland ecosystem enclosed by agricultural croplands and riparian vegetation. The presence of emergent vegetation and roosting trees offers ideal nesting and chick-rearing structures. This specific structural setup concentrates large avian flocks over narrow spatial boundaries, directly dictating localized nutrient influxes into the water and sediment. [1, 2, 3]

Ecological Significance of Guano at Udpuria Talab

1. Nutrient Loading and Geochemical Cycling

Fresh avian guano is highly concentrated in essential macro-nutrients, carrying roughly 11% to 16% Nitrogen (primarily uric acid), 8% to 12% Phosphoric Acid (phosphorus), and 2% to 3% Potash (potassium). [13,14,15]

- The Nutrient Vector Loop: Waterbirds that forage in surrounding agricultural crop fields but roost overnight at Udpuria Talab create a net import of terrestrial nutrients directly into the aquatic basin. [1, 2]
- Chemical Conversion: Herbivorous and piscivorous species breakdown complex organic elements, expelling them as bio-available nitrates (NO_3^-) and phosphates (PO_4^{3-}), which rapidly alter the limnological chemistry of the pond. [1, 2]

2. Fueling the Trophic Web (Primary Productivity)

The immediate biological outcome of guanotrophy is the upregulation of the wetland's primary productivity. [1]

- Plankton Blooms: The surge in localized nitrogen and phosphorus levels triggers rapid growth spikes in phytoplankton and micro-algae populations. [1]
- Invertebrate and Fish Stocks: These micro-algal blooms sustain diverse communities of zooplankton and aquatic macro-invertebrates. These, in turn, scale up the carrying capacity for local fish species, securing a self-sustaining food web that attracts tertiary predatory birds like herons, harriers, and ospreys. [1, 2, 3, 4]

3. Wetland Macrophyte and Riparian Growth

Guano drops collected under localized roosting trees act as a potent, natural slow-release fertilizer. This nutrient boost stimulates the growth of emergent macrophytes (like reeds and sedges) along the shoreline margins of Udpuria Talab. Healthy macrophyte layers provide crucial cover for breeding, cryptic nesting spots, and natural bio-filtration that stabilizes the shoreline against erosion. [1, 2, 3, 4]

4. The Delicate Balance: Risk of Eutrophication

While guano deposition is a healthy driver of baseline productivity, it carries ecological risks if the system is compromised. Because Udpuria Talab sits in proximity to agricultural zones, it suffers from additive pressures like synthetic chemical runoff. [1, 2, 3]

- Over-saturation: An extreme congregation of avifauna, combined with shrinking water levels during dry summer months, can push nutrient loads beyond the threshold, leading to guanotrophication. [16,17,18]
- Deoxygenation: Excessive organic loading can drop dissolved oxygen levels drastically, stressing fish stocks and increasing the threat of pathogenic outbreaks (like avian botulism or coliform multiplication). [1, 2]

Summary: Guano-Driven Ecosystem Mechanics

Ecosystem Metric	Direct Impact of Guano	Long-term Ecological Value
Water Chemistry	Elevates NO_3^- and PO_4^{3-} levels.	Shifts the trophic state to support complex life.
Primary Production	Stimulates phytoplankton and macrophyte growth.	Increases the biological carrying capacity of the wetland.
Secondary Production	Multiplies invertebrate and fish biomass.	Enriches foraging guilds for predatory avifauna.
Soil Quality	Deposits rich organic matter underneath roosting areas.	Enhances riparian vegetation structure and shoreline stability.

Aquatic bird guano is the accumulated excrement, eggshells, and remains of marine and coastal birds (such as cormorants, pelicans, boobies, and gannets). [1, 2, 3]

Key Characteristics & Composition

- Nutrient Richness: Bird guano is a highly potent organic fertilizer containing approximately 11% to 16% nitrogen, 8% to 12% phosphoric acid (phosphate), and 2% to 3% potash (potassium). [1, 2]
- Dietary Origin: Produced by fish-eating seabirds, whose marine diet concentrates high levels of nitrogen, phosphorus, and minerals in their waste. [1, 2]
- Dry Climate Formation: Accumulates in thick layers primarily in rainless, hot, and dry coastal regions and remote islands (such as the Chincha Islands off Peru or parts of Namibia) where the arid sun bakes the deposits into a dry, powdery form. [1, 2, 3]

Historical & Economic Significance

- Ancient Agriculture: Used for over 2,000 years by indigenous Andean societies—including the Chincha Kingdom and Incas—to supercharge maize (corn) production in desert soils. [1, 2, 3]

- The 19th-Century Guano Boom: Sparked a global trade and resource rush (including the U.S. Guano Islands Act of 1856 and the Guano Wars) before synthetic nitrogen fertilizers (Haber-Bosch process) largely replaced it in the 20th century. [1, 2, 3, 4]
- Modern Uses: Still harvested under strict, sustainable state management (such as in Peru) for organic and specialty farming. [1, 2]

CONCLUSION

Guano (accumulated bird or bat excrement) acts as a powerful nutrient and chemical driver in wetland ecosystems, influencing plant growth, water chemistry, and toxin accumulation. [1, 2]

Key Ecological Roles in Wetlands

- Nutrient Enrichment: Seabird and wading bird guano injects high levels of nitrogen, phosphorus, potassium, and organic matter into wetland soils and tree islands. This nutrient boost accelerates plant productivity and helps stabilize fragile wetland or coastal vegetation. [1, 2, 3]
- Mercury and Toxin Transport: Wading birds and seabirds can shuttle heavy metals—specifically toxic methylmercury (Hg)—into wetland food webs via their guano. In specific areas like the Florida Everglades, guano deposition rates for mercury have been measured at significantly higher rates than direct atmospheric fallout. [1]
- Trophic Base and Microbial Activity: Guano stimulates heterotrophic microbes, specialized bacteria, and fungi that break down organic waste, shaping localized aquatic and benthic food chains.
- Over-Enrichment Trade-offs: Excessive accumulation can lead to local phytotoxicity ("plant-burning"), severe odors, altered pH levels, and shifts in native aquatic plant and invertebrate compositions. [19,20]

REFERENCES

1. "Nauru: An Island Country Destroyed by Phosphate Mining". www.amusingplanet.com. Retrieved 24 October 2022.
2. Anuario de estadisticas del cobre y otros minerales [Yearbook: Copper and Other Mineral Statistics: 2004 2023] (Report). Comisión Chilena del Cobre. 2024.
3. "AAPFCO Publications". www.aapfco.org. Retrieved 24 October 2022.
4. "CDFA Bat Guano Labeling Guide" (PDF). www.cdfa.ca.gov/is/ffldrs/pdfs/BatGuano.pdf. Retrieved 28 October 2022.
5. "The Life Hidden Inside Caves: Ecological and Economic Importance of Bat Guano - Sakoui - 2020 - International Journal of Ecology - Wiley Online Library". 23 May 2025. doi:10.1155/2020/9872532. {{cite web}}: Missing or empty |url= (help)
6. Cushman 2013, p. 3.
7. Bongers, Jacob L.; Milton, Emily B. P.; Osborn, Jo; Drucker, Dorothée G.; Robinson, Joshua R.; Scaffidi, Beth K. (2026). "Seabirds shaped the expansion of pre-Inca society

in Peru". PLOS ONE. 21 (2) e0341263. Bibcode:2026PLoSO..2141263B. doi:10.1371/journal.pone.0341263. PMC 12893552. PMID 41671212.
8. Cushman 2013, p. 8.
9. Cushman 2013, p. 170.
10. Szpak, Paul; Millaire, Jean-Francois; White, Christine D.; Longstaffe, Fred J. (2012). "Influence of seabird guano and camelid dung fertilization on the nitrogen isotopic composition of field-grown maize (Zea mays)". Journal of Archaeological Science. 39 (12): 3721–3740. Bibcode:2012JArSc..39.3721S. doi:10.1016/j.jas.2012.06.035.
11. Santana-Sagredo, Francisca; et al. (February 2021). "'White gold' guano fertilizer drove agricultural intensification in the Atacama Desert from ad 1000". Nature Plants. 7 (2): 152–158. Bibcode:2021NatPl...7..152S. doi:10.1038/s41477-020-00835-4. ISSN 2055-0278. PMID 33495555. S2CID 231713231.
12. Schnug, Ewald; Jacobs, Frank; Stöven, Kirsten (5 November 2018). Guano: The White Gold of the Seabirds. IntechOpen. doi:10.5772/intechopen.79501. ISBN 978-1-78923-657-6. S2CID 133739523.
13. Cushman 2013, p. 26.
14. Cushman 2013, p. 29.
15. Miles, Wyndham D. (1961). "Sir Humphrey Davie, the Prince of Agricultural Chemists". Chymia. 7: 126–134. doi:10.2307/27757209. ISSN 0095-9367. JSTOR 27757209.
16. Cushman 2013, pp. 40–43.
17. Cushman 2013, p. 54.
18. Cushman 2013, p. 55.
19. Dwyer, Jim (10 June 2001). "June 3–9; The Root of a Famine". The New York Times. p. 2.
20. Cushman 2013, pp. 68–69.